

*Second part. K L 1857*


AN ESSAY ON

*The Physiology of Reproduction,*

*and*

*History of the Corpus Luteum.*

RESPECTFULLY SUBMITTED TO THE FACULTY OF THE



HOMOEOPATHIC MEDICAL COLLEGE  
OF PENNSYLVANIA,

*On the 28<sup>th</sup> day of January, Eighteen Hundred and  
Fifty-seven.*

*By E. W. Sanford.*

*Of Raynham, Bristol Co. Massachusetts.*

# The Physiology of Reproduction, and History of The Corpus Luteum.

Reproduction may be defined, as the act of a female being or animal, producing, or giving birth to another being or animal, constituted in all respects, like the parent.

In treating this subject, we shall confine our remarks to an exposition of the phenomena exhibited by the human female in the performance of the reproductive function.

In this department of Physiology, much may be said, yet little is known with positive accuracy.

Writers on obstetrics advance many opinions, and conjectures, yet comparatively few facts can be substantiated. Though many



have instituted laborious investigations, still the function is enveloped in mystery, and it is unlikely that human art will ever penetrate it.

What we do know, however, that two dissimilar systems operate harmoniously, - the one - female, - germiferous; the other - male, - possessed of spermatozoa, having vivifying qualities. The ability of these systems of genital organs, to perform, each its own proper function, is dependant upon many conditions.

### Action of the Male.

Procreative power seldom exists until the age of from fourteen to sixteen years.

Though prior to this time, the testicles may secrete a fluid,

yet it lacks an essential part,  
and therefore has no life giving  
property, — no power to fecundate  
a germ. At puberty, however, when  
the male, not unlike the lower ani-  
mals, receiving impressions through  
the senses of sight or touch, has  
connection with the female,  
the glans penis, in which are sen-  
sory nerves, is rendered acutely  
sensitive by friction against the  
rigid walls of the vagina, and  
an impression being conveyed  
to the Spinal Cord, reflex contrac-  
tion of the muscles of the Vesiculae  
Seminales, and of the Prostate  
gland, is produced; whereby the  
contents of these receptacles, mixed  
with, and diluting the testicular  
secretion, sui generis, forming



into the Urethra, from thence  
are forcibly ejaculated.

### Action of the Female.

We recognize the development of  
woman; we regard her as having  
arrived at a nubile age; that of  
puberty, by the performance of a  
seemingly necessary function—  
menstruation.

Menstruation is — Ovulation.

But we will define menstruation  
as a discharge of a sanguineous  
fluid from the Genitalia, occur-  
ring very regularly, once in 28  
or 30 days; sometimes more fre-  
quently. This act is dependent  
upon the existence of an uterus,  
with its appendages, the ovaries, wherein  
are lodged Graafian vesicles, en-  
closing ova, — germs.

The period of life at which this phenomenon is first observed, is usually at the close of the fifteenth year. The age, however, varies, in accordance with climate, diet, manner of living, and social relations. — The first discharge is of a sero-ranguineous character. The second assumes more the appearance of blood; and when the function is fully established, the menses present a dark, red color, bearing more resemblance to arterial, than to venous blood. — Where does this blood come from? Writers differ in their views respecting the source of the menstrual discharge; some considering it as a secretion of the mucous membrane lining



The uterus; others as an uterine hemorrhage. Analyses of the menses go to show that it is not dissimilar in its character to blood proper. The solid constituents of healthy blood are 210; Those of the menses 175. In the latter, blood corpuscles are found, and blood corpuscles are solids. Can solids be secreted? Fluids only are secreted. Therefore we may safely say that the menstrual discharge is an uterine hemorrhage, mingled with the mucus of the vagina. Appearing as an evidence of nubility, the 45<sup>th</sup> year usually brings with it the climacteric. The ability to bear children no longer exists. This uterine hemorrhage, or menstrual

flux continues from three to five days; and the amount of blood thus lost at each catamenial period, may be estimated at from four to six ounces.

Many women, however, have a much more copious flow of blood than this, without apparent detriment. If there are no ovaries; if there is no uterus, there can be no menstrual flow, nor, consequently, can she reproduce. With the loss of the ovaries only, but an imperfect sexual sense remains, often none at all.

But in the absence of the uterus, only, there may be sexual desire, but there can be no fecundation, - no conception. Many cases are recorded of women, who never had uteri,



and of necessity never ~~menstruated~~.  
Though this is a species of Amenorrhoea,  
it should not be mistaken for the  
Suppression; it is the Ennansio men-  
sim. Suppression of the menses  
can not be said to occur until  
after the full establishment of  
the function. Menstruation is  
Ovulation. The periodical he-  
matic discharge from the uterine  
vessels, is accompanied by the  
evolution and expulsion of an ovum,  
containing its ovule. —

Before proceeding, in detail, to  
speak of the separate phenomena  
of reproduction, we must devote  
some space to the Physiological  
consideration of certain organs,  
and their characteristics, all, par-  
ticularly concerned in the due per-

performance of the function. —

The Ovaria. Attached at each angle of the fundus of the uterus, at the distance of an inch and a half, lying behind the Fallopian tubes, are two bodies, of the size of an almond, enclosed in a duplicature of the peritoneum. Formerly they were called *Testes muliebres*; now known as ovaries, whose office it is to produce or prepare germs.

Covered by a peritoneal coat, the inner, or *tunica albuginea*, encloses, as in a shut box, a dense cellular tissue called *stroma*. In this stroma are imbedded numerous small vesicles or *buccae*, which, when freed from their attachment, appear globular, of the size of a pin's head. These spheroidal



10  
bodies are filled with a pellucid  
fluid, which, from its being coag-  
ulable by heat, is believed to be prin-  
cipally albuminous. It is in fact,  
vitelline matter, like the vitellus  
or yolk of an egg. These globules,  
first discovered by De Graaf, in 1672,  
are called Graafian vesicles, cells,  
follicles, or ova. In each ovary  
may be found 12 or 15 of these vesicles.

Previous to the age of puberty, how-  
ever, it is believed they do not exist;  
nor after the "critical period" of life.

Within these Graafian vesicles, or  
ovules, as Dr. Barry prefers to call them,  
are found numerous small granules;  
among which is an agglomerated  
mass, a cumulus of granules; and in  
the centre of this is the yolk ball,  
or ovum. Outside of the ovum or yolk.

ball, is a white, transparent circle, filled with oil globules, and vitellary corpuscles, called the zona pellucida. Within the ovum, near its periphery, is a clear, transparent, oval vesicle, in which, high microscopic power reveals a dark spot. The oval vesicle is the germinal vesicle, sometimes called the Purkinjean; and the dark spot is the germinal spot. It should have been stated above, that a Graafian vesicle has two coats; an outer, and inner coat. The inner one is called the membrana granulosa; and the granules forming the cumulus, outside the zona pellucida, are remains of the granular membrane. This membrana granulosa is thickest

in that part of the vesicle which is  
nearest the lining membrane of the  
ovary, - the tunica albuginea.  
Here a small elevation is formed, -  
an acromus, which Baer calls the  
proligerous disk (proles & zero). The  
germinal vesicle is  $\frac{1}{60}$ , and the  
germinal spot, the macula  
germinativa,  $\frac{1}{300}$  of a line in diam-  
eter. - Upon this point L. & C. writers  
now agree: - That at every menstru-  
ation, a mature ovum is expelled  
from its ovivac, - the Graafian ves-  
icle, and from the containing  
ovarium. Just before this rupture  
takes place, (the ovum now nearly  
matured,) the germinal spot moves  
towards the periphery of its vesicle;  
the germinal vesicle to that of the  
ovum; the ovum to that of the ovivac;



and the ovisac, to the circumference of the ovary. Thus when all the intervening walls are broken down, the seminal fluid may meet the germ, or ovulum; or, unstimulated by the sexual act, the extruded ovum falls into the peritoneal cavity, or as is more common, loses its vitality in a longer or shorter time, and then is destroyed, absorbed.

### The Corpus Luteum.

Upon the supposition that ova were never extruded except to be fecundated, writers have asserted that corpora lutea were the result of impregnation, and subsequent conception, alone. Now however it can no longer be denied, that a corpus luteum, at every catamenial period, is formed,

77

whether the extruded ovulum be fecundated or not. Every menstrual discharge proves the evolution and expulsion of an ovule; at what particular period, whether during, or at the termination of the flux, we do not know.

— For the sake of distinction, in which, though contrary to the belief of some writers, there is a difference, corpora lutea are spoken of as true and false; the former being those of pregnancy, and the latter of menstruation.

— Perhaps it may not be uninteresting to bring to mind the opinions of various eminent authors, concerning the corpora lutea.

William Hunter, and many Physiologists contemporaneous with him, disbelieving in the spontaneous evolution of ova, at

every Catamenial Period of the virgin  
considered the formation of the corpus  
luteum, as the accompaniment only  
of pregnancy; and necessarily con-  
nected with the act of impregnation.  
- Velpeau regards the corpus luteum  
as growing "after coitus." Dr. Gross  
says that false corpora lutea  
may be produced without coitus,  
as a consequence "of strong sexual  
excitement"; yet thinks the true cor-  
pus luteum a good evidence of preg-  
nancy. Mr. Haighton (1821) says  
that when corpora lutea are found,  
"they furnish incontestable proof  
that impregnation either does exist,  
or has preceded." Haller (1803) asserts  
the same. Montgomery (1831) says the  
growth of a corpus luteum takes place  
"on the occurrence of conception"



70

A host of authors, some even of recent date, have expressed their disbelief in the existence of corpora lutea in the virgin.

Dr. Robert Knox thinks "there is no distinctive character by which the corpus luteum (of impregnation) may be distinguished from the miniature corpus luteum"; or that formed by menstruation. Dr. G. Wharton Jones declares "it would be rash and unwarrantable for any one to pronounce, from the occurrence of a corpus luteum, in the ovary, that coitus had taken place." Bischoff (1847) was the first to announce the following theory:-

"That ova in mammalia, in the time of heat, no coitus taking place, are detached from the ovaries, enter the tube, and perish there; and that corpora lutea are found in the ovaries, just as though

cotion and fecundation has taken place". Dr. Dewees adopts the opinion of Sir. E. Home, that the corpora lutea exist previous to impregnation; "and they have no less a destiny to perform, than to furnish the ovum, and prepare it for impregnation."

M. Pouchet, admitting the existence of corpora lutea in the virgin, as well as in the mother, says there is no distinction between them, but that "they all have the same form, and the same structure." — Dr. John

C Dalton, Jr., however, declares "that the corpus luteum of pregnancy is different from the corpus luteum of menstruation; and that it may, under ordinary circumstances, be readily recognized and distinguished from it." (Vid essay, 1851). Adopting <sup>his opinion</sup>, we will

endeavour to furnish the proof of his assertion. As a medico-legal question, it may sometimes be necessary to inquire, - What are the characteristics of a corpus luteum of pregnancy? and what those of menstruation?

1<sup>st</sup>. What is the character of a corpus luteum of menstruation?

Recent investigations compel us to believe, that upon the accession of the menstrual discharge, - at puberty, an ovum, within its ovine, matured, protrudes from the surface of the ovary, bursts its walls, and becomes filled with blood. It not infrequently happens that this takes place in both ovaries at the same time; or that two Graafian vesicles are expelled from the same ovary. The quantity of blood thus effused,



is variable. Why it should be so, we can not tell, any more than we can assign a reason for the great difference existing in the size of ovaries.

On the extrusion of an ovum from the ovariole, the proligerous disk (page 12) at its base, and a portion of the granular membrane are expelled with it. At the same time, there is a deposition of a "fleshy, yellow looking substance," upon the proper membrane of the vesicle. This is regarded as an hypertrophy of the vesicular membrane, which, near the cicatrix is very thin. Meanwhile, the membrane is folding upon itself, presenting numerous convolutions, such as we meet with in the brain, to make the bulk proportionate with the cavity. As these changes progress, the effused blood becomes paler, in

"proportion to the time elapsed," since the rupture of the vesicle. By the third week from the last menstruation, the yellow body has attained its full size. Externally, its situation is indicated by a tumor, as large as a pistol bullet, soft and yielding under pressure. A coagulum fills the tumor, the yellow walls of which are  $\frac{1}{8}$  of an inch in thickness. The convoluted membrane, forming this wall, if spread out, would be much thinner. At this time, in the third week, the corpus luteum begins to retrograde, - to diminish in size. The coagulum, becoming still paler, is absorbed. From a sphere, the body is flattened. The retrogression continues till the sixth or eighth week from its first appearance. It then remains apparently stationary

till the eighth month, when it disappears.

This is the history of one. Meanwhile, many are in different stages of progression and retrogression, according to their various ages.

2<sup>nd</sup> ". What are the characteristic appearances of a corpus luteum of pregnancy?— The description of a corpus luteum of menstruation, i.e., following the expulsion of an ovum, of course involves no act of fecundation. Yet the development of a corpus luteum attendant upon pregnancy, in its commencement, is the same as above. But if the discharged ovum is impregnated, here commences the distinction. In the latter case, instead of attaining its full size in three weeks, and then becoming atrophied, and finally



disappearing, the corpus luteum of pregnancy "continues to develop itself;" increases in size, nor commences to retrograde till after parturition.

There is another distinction, - The process of vesicular protrusion ceases, for the time. Examinations of the corpus luteum of menstruation give evidence that its walls become much reduced in size, long before their yellow color fades away; that the coagulum retains its red appearance, even after the body is much diminished. In the case of a corpus luteum of pregnancy, these conditions are reversed. The yellow of the wall, and the red of the coagulum, seldom remain after the second month, while, meantime, the corpus luteum itself grows, - increases in size. At what period of

23  
uterus-gestation the yellow body is at its maximum, writers are not agreed. Dr. Carpenter states it as being between the third and sixth month. It is believed, that between these periods, the change, if any, is slight. - After parturition it rapidly diminishes, though it may be distinguished for many months. - In concluding this part of our essay, we will state the distinctions between corpora lutea of pregnancy, and those of menstruation, as given by Dr. Dalton, than whom, no one is more competent to decide. -

1<sup>st</sup>. It (i.e. the corpus luteum of pregnancy) arrives more slowly at its maximum of development, and afterwards remains for a long time as a very noticeable tumor, instead of under-



-going a process of rapid atrophy.

2<sup>nd</sup>. It retains a globular, or only slightly flattened form, and gives to the touch a sense of considerable resistance and solidity.

3<sup>rd</sup>. Internally, it has an appearance of advanced organization, which is wanting in the corpus luteum of menstruation.

4<sup>th</sup>. Its convoluted wall, particularly, attains a greater development, this portion remaining sometimes so much as  $\frac{7}{16}$  to  $\frac{1}{4}$  of an inch in thickness, while in the corpus luteum of menstruation it never exceeds  $\frac{1}{8}$ , and is almost always less than that. This difference in the thickness of the convoluted wall is one of the most important points of distinction. It will be much more striking when



viewed relatively to the size of the  
central coagulum.

5<sup>th</sup>. The color is not, by any means, so decided a yellow, but a more dusky, and indefinite hue.

6<sup>th</sup>. If the period of pregnancy is at all advanced, it is not found, like like the corpus luteum of menstruation, in company with unruptured vesicles in active process of development.

Vind Prize Essay. —

The Reproductive function may be comprised under five distinct heads, as follows:—

- 1<sup>st</sup>. Fecundation, or Impregnation.
- 2<sup>nd</sup>. Conception, or "Fixation" of the Germ.
- 3<sup>rd</sup>. Utero-gestation, or Pregnancy.
- 4<sup>th</sup>. Parturition, or Accouchement.
- 5<sup>th</sup>. Lactation, or Allaitement.

Perhaps it would have been well

the peace Generation, or the formation  
 of the Germ, at the head of the list;  
 But as we have spoken at great length  
 in this connection, relative to the over-  
 ry and its product, we will consider  
 1<sup>st</sup>. Fecundation or impregnation.  
 The terms fecundation and conception,  
 have been used by some authors,  
 synonymously. This cannot be.  
 Compelled by the mysterious ere-  
 stic influence, the Fallopian  
 tube approaches its ovary, grasps it  
 with its potaneous fimbriae, encloses  
 the ripened Graafian vesicle, presses  
 therefrom the mature ovulum, now  
 ready for the transit, which, entering  
 the tube, meets in its passage the  
 vitalizing sperm of the male;  
 becomes fecundated, - impregnated;  
 and then is conveyed to the uterine



cavity. This is Fecundation.  
But Fecundation is not Concep-  
tion. A fecundated ovule, unat-  
tached, may be lost in a sanguin-  
eous or mucous discharge, before  
conception can take place. But,  
remaining as it does, in a ma-  
jority of cases, within the womb,  
it becomes attached, - it has con-  
ceived. This is the theory. Mr. Aber-  
nethy declared "we know nothing  
of the phenomena of fecun-  
dation."

2<sup>nd</sup>. Conception (con & copio.)

Conception may be defined  
as the "fixation," (Keigs.) the attach-  
ment of an impregnated, - a fecun-  
dated germ or ovule, within its prop-  
er receptacle. The body of the uterus.  
A fecundated ovulum attaches



itself only to a mucous surface.  
If it therefore falls from the fimbriae,  
into the cavity of the pelvis, upon the per-  
itoneum, - a serous membrane, it is  
lost. Its only proper place is within  
the uterus. Conception, however, may  
take place elsewhere than in the womb,  
as in the Fallopian tube; in the ovarium;  
or in the substance of the walls of the  
womb. These are cases of extra-uter-  
ine pregnancy. - While the ova-  
rium, after fecundation, is in  
the Fallopian tube, the uterus is  
making preparations for its  
reception. A lining membrane  
is thrown around the entire cav-  
ity, closing up the tubular open-  
ings and the os uteri. This mem-  
brane is in two layers, between  
which, in the first months of pregnancy,

21  
a red colored fluid is found, called  
Hydroperione (ὕδωρ περιωρ.) This  
is finally absorbed, and the two  
layers are then fused together.

As the ovum enters the womb, this  
deciduous membrane is pushed be-  
fore it, reflecting or reflexing  
upon itself. To this part is given  
the name decidua reflexa. The  
decidua vera is the outer fold or  
layer of the whole membrane.

(Decidua, from decidere, to fall off)

The above is the Anterior Theory.  
M. Coste has proposed another, which  
Prof. Meigs partially adopts. (Viz Meigs)  
This deciduous membrane, then, is the  
outer covering of the embryo. The others  
also enclose it. - The Chorion (χορίον=  
the skin, from κορεν, to contain, to in-  
clude. This is internal to the decidua-



nous. Properly, it is the ocular membrane, formed while passing through the Fallopian tube, from feculent material, thrown out by the lining membrane; or perhaps from the zona pellucida (page 11) composed of oil-globules and granules, which were extruded with the ovulum.

The Amnion, (*braxeros*, a sheep, because first observed in that animal.) enclosing the liquor amnii, is the inner membrane of all, and comes in direct contact with the embryo. It is supposed to be produced by a fold of an external layer of the *membrana granulosa*, (page 11) spoken of as lining a Graafian vesicle, and surrounding the *zona pellucida*. It resembles the Chorion both in structure and appearance.



3<sup>rd</sup>. Pregnancy, or. Utero-gestation.

Pregnancy is the development of an embryo or fetus within the womb.

How soon this process commences after the fecundation of an ovule, and its subsequent conception, it is impossible to decide. During the first three months of utero gestation, the term embryo, is applied to the germ; after that time, or when it has become distinct in its outlines, we call it fetus. As pregnancy advances, there is an increase in the size of the womb, corresponding to the development of the embryo. In the virgin, the weight of the uterus is from 1 to  $\frac{1}{2}$  ounces; in length about 3 inches; and in breadth, 2 inches at the base, and 1 inch at the cervix. A wide departure from this standard attends the preg-

ren of lactation. At term, its gravid weight is from 1 to 1½ pounds; and instead of now being 3 inches in length it is 13 or 14. In this enormous growth of the uterus, its parietes do not become thinner; but in some instances are actually hypertrophied. The uterine arteries and veins elongate, attaining a greater proportional capacity. This is explained by the fact, that they are required to nourish not only the uterus itself, but to meet the urgent demands of the growing fetus. As gestation proceeds, the gravid uterus seeks more room. It rises from the pelvic into the abdominal cavity; often encroaching upon the intestines. The stomach now gives signs of sympathetic distress, evinced by nausea and vomiting. As physical signs of



It has been supposed that the presence of Kiesteine (from KUEIN = to be pregnant, & ΕΟΔΥΣ = a garment or pellice) in the urine, was an invariable diagnostic sign of pregnancy. - But Dr. E. F. Elliot Jr. of New York, in a recent paper, gives it as his belief, based on numerous observations, that this by no means can be relied on.



83  
pregnancy, we have great pro-  
trusion of the abdomen; though  
this is by no means always caused by a  
gravid womb. The umbilicus pro-  
trudes. Meanwhile, the menstrual  
discharge has ceased. By the fifth  
month, very convincing evidence of  
pregnancy is adduced, by a change  
in the position <sup>of the uterus,</sup> supposed to be a move-  
ment of the fetus, now vulgarly  
termed quickening. As a medico-legal  
point, the fetus is now recognized as  
possessing life, - is vivus, and  
to destroy it is a felony. Now,  
though prior to this time, there  
may be no movement of the fetus,  
it is not because there is no vi-  
tality; but from want of strength;  
and the movement, if there be any,  
is so feeble, that the mother is not conscious of it.

The ordinary term, or duration of utero-gestation, usually ranges between 9 calendar or solar months, (270 days), and 10 lunar months, (280 days.) Instances are recorded, however, of pregnancies continuing much longer than this; even to the 12<sup>th</sup> or 14<sup>th</sup> month. (Meigs.) Again, the child has been born in less than the usual time. But 280 days may be stated as the usual term.

4<sup>th</sup>. Parturition or Accouchement. Parturition, termed Tocology, (tokos birth, & logos) by Dr. Ryan, is a function, which consists in the expulsion of the contents of the gravid uterus. It is Labor; and may be natural, or unnatural. Happily a vast majority of cases belongs to the former condition. Different

writers on Obstetrics, divide parturition or labor, into various stages, or periods. The following by Denman is considered the best arrangement.

There are three stages or periods.

1<sup>st</sup> Stage, Commencing with the accession of labor pains, extends to the dilatation of the os uteri, and the consequent discharge of the liquor amnii.

2<sup>nd</sup> Stage, Commencing with the discharge of the "waters", continues until the birth of the child.

3<sup>rd</sup> Stage, Commencing with the delivery of the child, continues till the expulsion of the placenta, and membranes, - the secundines, or "after birth."

The first stage lasts from one, to



six hours. The second from one to three or four hours; and the third usually from fifteen to twenty minutes. Each of these periods is characterized by its own peculiar phenomena.

5<sup>th</sup>. Lactation, or Alactement.

The mammary glands exist in both sexes, and are situated immediately over the great pectoral muscles; a layer of superficial fascia only intervening. In structure, this gland is like that of any other; composed of lobes and lobules, - acini, connected by a fibrous or fascial tissue. Each of these glandules gives out a tube, or lactiferous vessel, which, joining with others, form a communication to the

external surface. There are from fifteen to twenty of these ducts, or galactophorous tubes, in each gland. As they approach nearer the surface, they become larger, forming reservoirs, which, at parturition, hold in reserve a "draught,"—a milky secretion, to supply the immediate wants of the infant. Up to the age of puberty, the mammary gland of the female, is in no respect different from that of the male. At that time, however, or as the period approaches, the gland of the female rapidly enlarges; and the lactiferous ducts become capable of injection. External to the gland itself, the part increases in size; caused by a deposition of masses of adipose matter. At the age of

minute or twenty, the gland  
 has attained its maximum of  
 development; the mamma is  
 fully formed, capable of per-  
 forming that function, which  
 necessarily devolves upon it at  
 pregnancy. — In the centre of  
 the breast, is a conoidal eleva-  
 tion, — the nipple. Around this  
 extends a circle, which, in the  
 virgin assumes a rosy tint; but  
 as pregnancy advances, this circle,  
 — the areola, becomes of a darker  
 color, wider in its outline, and  
 is of a rugous character, contain-  
 ing a number of sebaceous follic-  
 les or glands. — In man, the  
 mammary gland differs only  
 in size from that of the woman.  
 Instances are recorded of men,



In the mammary secretion previous to parturition, albumen is found to be the principal ingredient; with scarcely a trace of sugar. Gradually Casein takes the place of the Albumen, and at the same time, sugar and fat appear. After delivery, the secretion called Colostrum, (vid 40<sup>th</sup> page) is found. This is thicker than true milk, having a light yellow color, with no peculiar odor. It is alkaline in its reaction. The taste is very sweet. Analysis shows it to be much richer in butter and saccharum lactis, than ordinary milk.

whose glands and mammae became largely developed, in consequence of permitting a child to suck the rudimentary nipple. This anomaly has been observed in some of the lower animals.

— The secretion of the mammary glands, — milk, is composed of water, holding in solution, sugar, salts, and an albuminous compound called caseine.

This caseine contains oleine, which decomposes the proximate principle of milk, — butter, making it rancid. — Woman's milk contains more sugar and less caseine, than cow's milk; and in these respects resembles ass's milk.

*Saccharum Lactis* is obtained by evaporating the whey of milk,



It is stated by French Physiologists,  
that during the period of suckling  
the infant, Glucose, a kind of  
sugar, is found in the urine of nur-  
sing women. Under these circum-  
stances, there is no evidence of  
the existence of Diabetes Mellitus.  
This sugar exists in the propor-  
tion of 6 parts in the 1000.



and allowing the syrup thus formed, to crystallize. ( $C^{24}O^{24}H^{24}$ )

At parturition, the first secretion of milk is known by the name of Colostrum (Kolos = food.) This colostrum has a purgative effect upon the child, causing an evacuation of the meconium, ( $\mu\eta\kappa\omicron\varsigma$  = poppy, an accumulation of fecal matter in the intestines.) At the expiration of twelve months, the colostric character of the mother's milk returns, and to nurse the child any longer, is highly injurious. Recourse must be had to other food. Dr. Rees states, that milk, in its composition, resembles blood, more than any other fluid.

TELOR.

E.M.P.